

Preparing for Foot and Ankle Surgery

A Patient's Guide

This booklet is intended to give you an idea of what is involved in undertaking foot and ankle surgery. It provides information to help your decision, with your surgeon, on whether surgery is the right treatment for **you**.

When your consultant offers you an operation, their decision is based upon their experience and expertise and numerous factors related to you.

You should also be aware, however, that:

- **Surgery should be considered to be the final option**

Foot and ankle problems can be treated in many ways without having an operation. These include custom-made insoles and/or shoes, physiotherapy and exercises, medicines and injections, weight loss or a change in activity level. Surgery may be offered if these treatments are unlikely to help or have failed to improve your symptoms.

- **Surgery aims to improve your function**

This occurs by reducing pain and helping you to walk more efficiently. ***Surgery is not performed for cosmetic reasons.*** Satisfaction with an operation to improve the appearance of your foot is unpredictable and the risks of surgery outweigh the benefits. Your surgeon will not offer you an operation for a purely cosmetic problem

This page is intentionally left blank.

- **You must have realistic expectations of surgery**
We perform surgery to enable patients to cope with day to day activity. However, this may involve changing the way the foot or ankle works forever. The foot does not always respond kindly to injury or surgery and can take many months to recover. It is **unusual** for the foot to be perfect when it comes out of bandaging or plaster and, in many cases, it may be 12-18 months before the final outcome is appreciated.
- **You must take control of your treatment**
You will need to follow all instructions given for your operation. Failure to do so can have detrimental long-term consequences. For example, if your foot is not elevated after surgery, swelling can quickly occur, leading to pain and increased risk of delayed wound healing and infection. Successful results require a contribution from the patient, as well as the surgeon and the hospital team.
- **The success rates of surgery are generally high**
For some problems, however, there may be no operation available to reliably improve your symptoms; in these cases surgery may carry unnecessary risks. Rarely, in cases of extreme suffering, you may be offered surgery even when the chance of success is poor. However, this would only happen after thorough discussion and your acceptance of this increased risk.
- **All surgery carries risks**
Your surgeon will talk you through the operation and the most relevant complications that could potentially occur.

You should think about these and ask for clarification if you are uncertain about them. As surgeons, we aim to minimise complications as much as possible, but they still occur. This booklet includes a list of complications that could happen to you. Fortunately, they are not common but, *if you cannot accept the risk, do not have the operation.*

Before Your Surgery

You will be asked about your general health and smoking status when you are put forward for surgery. A further pre-operative clinic check-up is also required in order for you to undergo your operation. This can highlight any health problems that may give you an increased risk of complications. If these can be remedied or improved upon before surgery, then it may be appropriate to postpone your operation.

- **Smoking**

If you are a smoker, you should seriously consider stopping completely before having a general anaesthetic and surgery. Smokers have more complications after surgery with a much higher risk of wound breakdown, wound infection, deep vein thrombosis (DVT) and failure of bony operations. The risk of complications and failure is so high after certain operations that your surgeon may be reluctant to perform it without evidence of you stopping smoking. If you wish to give up smoking, the hospital or your GP can offer you help and advice on stopping smoking.

Don't hesitate to discuss with your surgeon any questions you may still have after reading this booklet.

Internet sources of patient information you may find useful:

- www.arthritisresearchuk.org
(<http://www.arthritisresearchuk.org/arthritis-information/surgery/foot-and-ankle-surgery>)
- www.bofas.org.uk
- www.myankle.co.uk

Contacts

Tel: 01325 380100 (DMH) or 0191 3332333 (UHND) and ask for your consultant's secretary

Foot & Ankle Consultants at County Durham and Darlington Foundation Trust:

Mr Thomas Barwick
Mr Graham Chuter
Mr Andrew Gower

Production date: September 2015
Review date: September 2018
Responsibility for review: Mr T Barwick (orthopaedics DMH)
Leaflet reference: PIL/CG/0249
Version: 2.0

- Complex regional pain syndrome (unpredictable pain, swelling, stiffness and circulation changes)

Your surgeon will discuss any complications that are specific to your procedure.

General complications

- Heart attack
- Stroke or mini-stroke (TIA)
- Blood clots in leg veins (deep vein thrombosis) or lung (pulmonary embolism)
- Stomach/bowel ulceration (in response to 'stress' of surgery)
- Chest infection
- Allergic reaction to drugs or blood transfusion
- Temporary worsening of diabetes
- Protecting the foot after surgery can put additional strain on other joints e.g. hip and back

Some of these complications can be fatal but, fortunately, this is extremely rare.

Although this is a fairly frightening list you should remember that planned, modern surgery is very safe compared with many activities we all encounter every day.

Finally, we hope this leaflet assists your decision making when it comes to your treatment and helps to separate the truth from fiction when it comes to foot and ankle surgery.

- **Oral Contraceptive Pill**

The risk of Deep Vein Thrombosis (a blood clot in the leg) and pulmonary embolism (a blood clot in the lung) is higher in women taking oestrogen-containing oral contraceptive pills (even low dose). You should stop the pill a minimum of 4 weeks prior to elective surgery, and preferably 6 weeks before. If you do not know whether your pill contains oestrogen, ask your doctor or the clinic that prescribed it. You should also see them to get advice on preventing pregnancy whilst off the pill.

- **Hormone Replacement Therapy (HRT)**

HRT contains lower levels of oestrogen than the oral contraceptive but may still increase the risk of DVT. It is recommended that you stop HRT for two weeks prior to your operation date. If, however, the effects of stopping are too unpleasant, you may restart it, but we will then prescribe medication to help reduce the risk of DVT. These will start on the day of surgery and continue until you are fully weight-bearing.

- **Podiatry treatment**

If you are having chiropody or podiatry treatment, it is important that you tell your podiatrist that you will be having surgery. Your last treatment should be one week before the date of surgery at the latest. Do not resume any treatment after surgery without consulting your surgeon.

- **Athlete's Foot**

If you suffer with 'athlete's foot', you should try to have this cleared up before any surgery. It is acceptable to use

antifungal treatment regularly but do not let your foot get too moist from excess use. Treat your socks and footwear with antifungal powder to prevent re-infection. Ask your GP or pharmacist for advice if you need help treating this common problem.

- **Other Medicines and Allergies**

Some medicines cause problems with anaesthesia and surgery. You should inform the Pre-operative assessment team of all medication and non-prescription medication that you take. You may be allergic to some hospital drugs that we use routinely; please inform us of any allergies you may have.

- **Extra help you will need after surgery**

Plan to rest; you may be far less mobile than normal for some time. Ensure you have friends or family to help as necessary. If you live alone, you will need to consider help with certain tasks like shopping and preparing meals. Will you be able to negotiate your house/ stairs with crutches/ sticks? Inform the clinic as soon as possible about your needs and the necessary help can be arranged. You may need an overnight stay on the ward in certain cases.

Before Admission

At the **Pre-operative assessment clinic** you will be asked many questions about your health. Tests will also be arranged as appropriate but typically include blood tests, heart tracings (ECG) and X-rays. Occasionally, the clinic highlights health problems that require further specialist tests and/or treatment to be carried

increases in activity without pain or swelling. This may be several months; your physiotherapist will help to supervise this.

The risks of foot and ankle surgery

Your surgeon will explain some of the risks that are involved with your surgery. Occasionally, even operations with predictably excellent results can result in a poor outcome when a complication occurs. We cannot always predict which patients will get a complication, but we can look for the conditions that may increase the chances of a complication occurring, such as diabetes or smoking. *It is for you, the patient, to decide what level of risk you are prepared to accept.* Your surgeon will use their knowledge of your risk factors, the surgical procedure and their experience to help you make an informed choice as to whether to proceed with surgery.

Complications related to the operation site

- Bleeding, bruising, swelling (long-term)
- Infection and wound breakdown
- Damage to blood vessels that may require surgical repair or result in poor circulation that may lead to loss of tissue or toes or the foot (amputation)
- Persistent tenderness or sensitivity near scars; ugly or raised scars
- Damage to major nerves leading to absent, abnormal or painful sensations or local weakness or paralysis of the foot or ankle

several days; your surgeon will determine when your wound is first re-dressed and/or inspected. This may be within the first week or at two weeks depending on the operation and the surgeon's clinic times. Plaster casts must remain dry at all times.

- **Weight-bearing**

Your operation may involve placement of metal ware and repair of soft tissues that cannot tolerate 'full weight-bearing' initially. Therefore, after your operation you might be instructed to be 'non weight-bearing'. Other forms of weight-bearing include 'heel weight-bearing', 'toe-touch weight-bearing' and 'partial weight-bearing'. These will be fully explained to you by the staff after your operation and must be strictly followed to prevent failure of surgery. Often we supply a special form of post-operative shoe to take the pressure off the site of surgery e.g. bunion surgery.

- **Driving**

You should not drive until you can wear a normal shoe and can perform an emergency stop with confidence e.g. stamping your foot gently on the ground. The responsibility to return to driving safely lies with the patient (contact the DVLA/ see DVLA website). Inform your insurance company about your operation after returning to driving.

- **Sport**

Discuss this with your surgeon before returning to activity. Non weight-bearing activities (such as swimming) may be tolerated soon after wounds have fully healed. Contact sports may be resumed only after controlled, gradual

out, necessitating a postponement of surgery. Although this causes frustration for everybody it cannot be avoided and is clearly in your long term interest.

Please ensure your foot hygiene is as good as possible on the days leading up to surgery. If you have difficulty with addressing your foot care, please arrange a timely visit to your chiropodist at least a week prior to surgery.

On the Day of Admission

Usually you will be admitted on the day of surgery to the Surgical Admission Unit (SAU) or equivalent, depending on whether you are attending Darlington Memorial Hospital, the University Hospital of North Durham, or Bishop Auckland Hospital. If you have difficulty arriving early in the morning, on the day of surgery, then we need to know in good time to prevent a delay to the start of the list.

Some patients with certain health problems may need to be admitted the day before surgery. If this is necessary, this will be discussed at the Pre-operative assessment clinic.

The Operation

Your surgeon and anaesthetist will see you before the operation. The surgeon will answer any further questions you may have, take verbal and written consent for the procedure (if not already given) and mark your leg. The anaesthetist will discuss their role and the options specific to your surgery; this includes general and/or local anaesthetic (numbing injections) as well as post-operative pain control. Local anaesthetic blocks around the foot and ankle have greatly improved the experience of patients in

recent years and may last for up to 24 hours. You are advised, however, to start regular pain killers well before it wears off.

At the time of surgery you will leave the ward and be taken to a 'holding bay' in the theatre suite for safety checks. Staff will then take you to the anaesthetic room, next to the operating theatre.

After the operation, your foot and ankle will either be bandaged or in a plaster cast. You will wake up in the 'recovery room' before being returned to the ward once you are awake and comfortable.

After the Operation

When in bed, you will be instructed to keep your leg elevated before attempting to get up. The nurses and physiotherapy staff will assist you when you are ready to mobilise on the ward. They will also provide further advice for when you are at home, especially if you still have a numb foot from the anaesthetic. You will not be discharged until you are safe to mobilise in the manner the surgeon intended.

Increasingly, foot and ankle surgical patients are discharged home on the same day of surgery. This depends upon the patient, the type of surgery or anaesthetic undertaken, and the surgeon's preferences. You should be informed at your Pre-operative assessment clinic so that you know what to bring with you on the day of surgery. *Please organise transport home as you may not be allowed to drive.*

When you get home

- **Elevate Your Foot**

This is vital to reduce swelling and to speed the healing process. For the first couple of weeks after surgery, keep your foot up for 50 minutes every hour, ideally at the level of your heart. As a guide, if you are not preparing meals, bathing or toileting, your foot should be elevated. Please do it until your wound has been checked in clinic and is healing as expected. Excessive swelling may also be reduced by applying ice (or a bag of frozen peas) wrapped in tea towels applied directly on to the dressings.

Reducing the swelling will reduce the pain!

- **Check Your Foot**

Observe the colour of the foot or tips of the toes. Bruising is normal, as is some oozing of the wound(s). However, blue or white toes, worsening pins and needles, or massive swelling needs urgent attention by medical staff. You should ring the ward where you had your operation, contact your GP or attend your nearest A&E department for an urgent assessment.

Other aspects of your care include:

- **Pain relief**

Take pain killers regularly (as prescribed); you may need them for several weeks after surgery (very variable). Discuss with your GP if you encounter any side effects.

- **Dressings and wound care**

Keep your bandages clean and dry. *Do not tamper with the dressings.* The initial dressings are usually left on